

Performance Audit on the Implementation of the National Agenda Plan for Particulate Matter (PM2.5) Mitigation

Preventing and Reducing Open Burning in Sugarcane Cultivation Areas








Reason and Importance


- Thailand has long faced the problem of particulate matter (PM) pollution.
- Since 2019, the Thai government has launched the National Agenda Plan for Particulate Matter (PM) mitigation.
- One major source is open burning in sugarcane fields, commonly burned before harvest to reduce cost.
- The government has allocated over 25 billion baht to phase out burnt sugarcane by the 2023/2024 crop year.

Source of Data

Analysis of Particulate Matter (PM 2.5) and Fire Hot Spot Situation

- PM2.5 data, estimated from Aerosol Optical Depth (AOD) from 2000 to 2024 were analyzed using the Google Earth Engine (GEE) platform, which provides satellite-derived datasets.
- PM2.5 data, Ground-based air quality data were sourced from the Pollution Control Department (PCD) monitoring stations.
- Fire Hotspot data, indicating thermal anomalies, were collected using VIIRS sensors onboard JPSS NOAA-20 (J1), JPSS NOAA-21 (J2), and SUOMI-NPP satellites, supported by the Fire Information for Resource Management System (FIRMS) operated by NASA.





Source of Data

Analysis of Burning Reduction in Sugarcane Cultivation

- Land use maps at 1:25,000 scale, provided by the Land Use Policy and Planning Division, Land Development Department
- Farmer's sugarcane delivery data and official sugar production reports, from crop years 2012/2013 to 2023/2024
- Payment records under the financial support program for farmers delivering fresh (non-burnt) sugarcane, from crop years 2019/2020 to 2021/2022







Data analysis tool














```
Jupyter DA Stack And Direction19.12.24-report... Last Checkpoint: Last Thursday at 9:59 AM (autosaved)
File Edit View Insert Cell Kernel Widgets Help
    In [1]: import pandas as pd
             import numpy as np
from sklearn.model_selection import StratifiedKFold
             from sklearn.ensemble import RandomForestClassifier, GradientBoostingClassifier, VotingClassifier,
                 StackingClassifier
             from sklearn.linear_model import LogisticRegression
             from xgboost import XGBClassifier
             from lightgbm import LGBMClassifier
             from catboost import CatBoostClassifier
             from sklearn.metrics import accuracy_score, classification_report, confusion_matrix, roc_auc_score,
                precision_score, recall_score, f1_score
             # 1. โหลดข้อมูล
             data_path = r"sugarcane_data.csv"
             df = pd.read_csv(data_path)
             # 2. กำหนดด้วแปรดันและด้วแปรเป้าหมาย
             X = df.drop(columns=['target1_0', 'no_cane_year_target'])
             # 3. จัดการค่าที่เป็น NaN
             X = X.fillna(X.median())
             # 4. ตรวจสอบ Outliers และตัดขอบบน-ขอบล่าง 1%
             for col in X.columns:
                lower_bound = X[col].quantile(0.01)
                 upper_bound = X[col].quantile(0.99)
                X = X[(X[col] \rightarrow lower_bound) & (X[col] \leftarrow upper_bound)]
             rf_clf = RandomForestClassifier(n_estimators=300, random_state=42)
             gb_clf = GradientBoostingClassifier(n_estimators=300, random_state=42)
xgb_clf = XGBClassifier(use_label_encoder=False, eval_metric='logloss', random_state=42)
             lgb_clf = LGBMClassifier(n_estimators=300, random_state=42)
             cat_clf = CatBoostClassifier(silent=True, random_state=42)
             lr_clf = LogisticRegression(max_iter=1000, random_state=42)
             voting_clf = VotingClassifier(estimators=[
                ('rf', rf_clf),
('gb', gb_clf),
                 ('xgb', xgb_clf),
                 ('lgb', lgb_clf),
                 ('cat', cat_clf)
             ], voting='soft')
```


Data analysis tool

Google Earth Engine

javascript









Data Analytics Results


- The Trend of PM2.5 and burning (fire hot spot) problems
- The situation of sugarcane burning and the behavior of farmers after subsidy receiving
 - Apply the Churn Analysis to examine individual farmer's behavior after receiving supports, the proportion of burnt cane continues to decrease?
- Key factors for reducing PM2.5 (burning) in sugarcane cultivation areas
 - Machine Learning (Classification Supervisor)
 - Analysis of Variance (ANOVA)




Concentration of fine particulate matter (PM2.5 level)



Estimated from Aerosol Optical Depth data from 2000 - 2024 (25 yrs)







Concentration of fine particulate matter (PM2.5 level)



Monthly PM2.5 levels and trends in 2000 to 2024 (25 Years)









March, April, May, and October




Fire Hotspot Situation in Thailand (2019-2024)






Hotspot Distribution across Land Use Categories Flourish*




Fire Hotspot Situation in Thailand (2019-2024)



Monthly Fire Hotspot Situation in Thailand (2019-2024)









Fire Hotspot Situation in Thailand (2019-2024)





The Relationship Between Hotspots and PM 2.5 (2020–2024)



Overview Key Findings

The government's actions to mitigate PM2.5 pollution have shown limited effectiveness. In other words, the measures implemented so far have proven insufficient to resolve the issue.








•• Focus on: Preventing and reducing Open Burning in Sugarcane Cultivation Areas






- **Measures Implemented**
- Financial Support for Fresh-Cut Sugarcane
 - 2019/2020 : 92 baht/ton
 - 2020/2021 2022/2023: 120 baht/ton
- Interest Subsidy for buying harvesters
- Lend leaf-stripping machines

- Penalty Deduction for Burnt Cane
 - 30 baht/ton
- Collaboration request to sugar factory

Sugarcane Delivery to Mills by Farmers

Target Proportion of Burnt Sugarcane (%)









Proportion of burnt sugarcane has exceeded the target in every production year.

Open Burning and Hotspot in Sugarcane Cultivation Areas






Number of Hotspots (points)




Sugar Mill : Burnt Cane Proportion in 2019/2020 - 2022/2023

The sugar mill that process a large proportion of burnt cane are mostly located in the northeastern, eastern, and central regions.



Daily Burnt Cane, Hotspots, Burnt Ratio, and PM2.5 - 2022/2023



Burnt Sugarcane Volume (Million Tons)

Number of Hotspots in Sugarcane Areas

Average PM2.5 in Sugarcane-Growing Provinces (µg/m³)

Percentage of Burnt Cane Delivered to Mills (%)

Pearson Correlation Coefficient

- PM2.5 ↔ Number of hotspots: r = 0.634
- PM2.5 ↔ Percentage of burnt cane: r = 0.519
- Number of hotspots ↔ Percentage of burnt cane: r = 0.441

Government Measures to Promote Fresh-Cut Sugarcane

in the following crop year

Cash subsidy (direct payment)

Crop 2021/2022








farmers delivered more burnt cane





8,110.96 million baht

114,890 farmers

Low-interest loans for farmers

Year 2018-2023








farmers delivered burnt cane





Leaf-stripping machines provided for use/loan


Crop 2022/2023





farmers delivered more burnt cane






Burnt Cane Penalty and Farmer Behavior



Leaf Cane and residue management Post-harvest


From field surveys






Post-harvest burning

Pre-harvest burning



Key Factors Influencing the Success in Reducing PM2.5 in Sugarcane Areas

The XGBoost model achieved the highest predictive performance, with Accuracy of 0.8242



Feature importance analysis

- Number of sugarcane harvesters available in the area
- Proportion of fresh cane harvested by machine (%)
- Period of cane-receiving at sugar mills



Policy Recommendations

- Expand Scope Beyond Sugarcane
 - Include other major agricultural sectors (e.g., rice, maize, cassava) under the anti-burning initiatives.
 - Burning in natural forests and newly reforested zones
- Shift from direct subsidy only toward area-based problem solving, like land issues or lack of machines
- Increase penalties for burnt sugarcane
- Enforce stricter sugar mill policies against accepting burnt cane.
- Support using machines for fresh cane cutting
 - Make fields suitable for machine use
 - Offer incentives to entrepreneurs and factory investing in machine harvesting.
 - Government, farmers, and sugar mills should plan together for planting and harvesting.

BENEFITS Applying Big Data Analytics in Auditing

Enhance Accuracy and Precision

- Analyze large datasets to reduce errors from traditional sampling methods
- Use in-depth data analysis to spot hidden patterns and to discover trend

The Improve Operational Efficiency

- Reduce the time to process and analyze data
- Use visual tools to help teams clearly understand and communicate findings.





Recommended next steps

- Strengthen data analytics capacity for auditors at all levels—through hands-on training, role-specific upskilling, and real-case practice.
- Promote a data-driven audit culture that enables deeper analysis and evidence-based conclusions.
- Develop a Modern Audit System with integrated tools and a Data Lab that connects to external agency databases.
- Provide structured support through Data Support Teams and Data Analysts who assist auditors throughout the audit process.

